arXiv:math-ph/0502048v1 25 Feb 2005

Group classification of systems of non-linear reaction-diffusion equations with general diffusion matrix. III. Triangular diffusion matrix

A. G. Nikitin
Institute of Mathematics of Nat.Acad. Sci of Ukraine, 4 Tereshchenkivska str.
01601 Kyiv, Ukraine

Absrtract. Group classification of systems of two coupled nonlinear reaction-diffusion equation with a general diffusion matrix started in papers [1], [2] is completed in present paper where all non-equivalent equations with triangular diffusion matrix are classified. In addition, symmetries of diffusion systems with nilpotent diffusion matrix and additional first order derivative terms are described.

1 Introduction

Group classification can be considered as one of basic elements of qualitative analysis of entire classes of differential equations. Such classification makes it possible to select the basic non-equivalent equations in the considered class of them. In addition, it opens the way to application of powerful group-theoretical tools for searching for conservation laws, construction of exact solutions, group generation of families of solutions starting with known ones, etc., etc.

In the present paper we complete the group classification of systems of reaction-diffusion equations with general diffusion matrix performed in papers [1], [2]. These equations can be written in the following form:

$$\begin{array}{l} u_t - \Delta(A^{11}u + A^{12}v) = f^1(u,v), \\ v_t - \Delta(A^{21}u + A^{22}v) = f^2(u,v) \end{array} \qquad (1)$$

where u and v are function of $t, x_1, x_2, \ldots, x_m$, A^{11}, A^{12}, A^{21} and A^{22} are real constants and Δ is the Laplace operator in R^m.

Equations (1) form the grounds for a great many models of mathematical physics, biology, chemistry, etc., which makes their group classification to be especially relevant. We will not discuss here the history of investigation of symmetries of systems (1) which can be found in [1], [2].

Up to linear transformation of dependent variables there exist three *a priori* non-equivalent versions of equations (1), corresponding to a diagonal, triangular and square diffusion matrix. The equations with diagonal and square diffusion matrix have been studied in papers [1] and [2] correspondingly were the complete group classification of such equations is carried out.

Here we consider the remaining case when the diffusion matrix is triangular and system (1) is reduced to the following form:

$$\begin{aligned} &u_t - a\Delta u = f^1(u,v),\\ &v_t - \Delta u - a\Delta v = f^2(u,v)\end{aligned} \tag{2}$$

where a is a real constant.

In the case when $a = 0$ the diffusion matrix is nilpotent and it is naturally to generalize (2) to the following system

$$\begin{aligned} &u_t - p_\mu v_{x_\mu} = f^1(u_1,u_2),\\ &v_t - \Delta u = f^2(u_1,u_2)\end{aligned} \tag{3}$$

were (and in the following text) summation from 1 to m is imposed over the repeated Latin indices, p_μ are arbitrary constants. Using linear transformations of independent variables we reduce values of p_μ to

$$p_1 = p_2 = \cdots = p_{m-1} = 0, \;\; p_m = p \tag{4}$$

where $p = \sqrt{p_1^2 + p_2^2 + \cdots + p_m^2}$.

We notice that in addition to possible application in mathematical biology equations (3) can serve as a potential equation for the nonlinear D'alembert equation.

2 Symmetries and determining equations

To describe symmetries of equations (2) and (3) with respect to continuous groups of transformations we use the Lie infinitesimal approach. Using the standard Lie algorithm we can find the determining equations for coefficient functions $\eta,\ \xi_a,\ \pi^b$ of generator X of the symmetry group:

$$X = \eta\partial_t + \xi^\nu\partial_{x_\nu} - \pi^1\partial_u - \pi^2\partial_v \tag{5}$$

The first set of determining equations describes dependence of η, ξ^ν and π^b on u and v:

$$\eta_u = \eta_v = 0, \;\; \xi^\nu_u = \xi^\nu_v = 0, \;\;\; \pi^a_{uu} = \pi^a_{uv} = \pi^a_{vv} = 0. \tag{6}$$

So from (6) η and ξ^ν are functions of t and x_μ, and

$$\pi^a = N^{a1}u + N^{a2}v + B^a, \quad a = 1,2 \tag{7}$$

where N^{ab}, B^a are functions of t and x_ν only. The remaining determining equations in the case $a \neq 0$ are [1]:

$$2A\xi^\nu_{x_\mu} = -\delta^{\mu\nu}(\eta_t A + [A,N]), \qquad \eta_{x_\nu t} = 0, \tag{8}$$

$$\begin{array}{l} \xi^\nu_t - 2aN^{11}_{x_\nu} - a\Delta\xi^\nu = 0, \quad N^{12} = 0,\ N^{21}_{x_\nu} = -aN^{11}_{x_\nu}, \\ \eta_t f^k + N^{kb} f^b + (N^{kb}_t - \Delta A^{ks} N^{sb}) u_b + B^k_t - \Delta A^{kc} B^c \\ = (B^a + N^{ab} u_b) f^k_{u_a}. \end{array} \tag{9}$$

Here N and A are matrices whose elements are N^{ab} and A^{ab}, δ^{ab} is the Kronecker symbol, and we use the temporary notation $u = u_1, v = u_2$.

Using (7)–(9) we find the general form of symmetry (5) admitted by equation (2):

$$\begin{array}{l} X = \Psi^{\mu\nu} x_\mu \partial_{x_\nu} + \nu\partial_t + \rho_\mu \partial_{x_\mu} + \lambda K + \sigma_\mu G_\mu + \omega_\mu \hat{G}_\mu + \mu D \\ -C^1(u\partial_u + v\partial_v) - C^2 u\partial_v - B^1\partial_u - B^2\partial_v \end{array} \tag{10}$$

where the Greek letters denote arbitrary constants, $B^1,\ B^2$ are functions of t, x, and $C^1,\ C^2$ are functions of t,

$$\begin{array}{l} K = 2t(t\partial_t + x_\mu \partial_{x_\mu}) - \frac{x^2}{2}\left(\frac{1}{a}(u\partial_u + v\partial_v) - u\partial_v\right) - tm(u\partial_u + v\partial_v), \\ G_\mu = t\partial_{x_\mu} + \frac{1}{2} x_\mu \left(\frac{1}{a}(u\partial_u + v\partial_v) - \frac{1}{a^2} u\partial_v\right), \\ \hat{G}_\mu = e^{\gamma t}\left(\partial_{x_\mu} + \frac{1}{2}\gamma x_\mu \left(\frac{1}{a}(u\partial_u + v\partial_v) - \frac{1}{a^2} u\partial_v\right)\right), \\ D = t\partial_t + \frac{1}{2} x_\mu \partial_{x_\mu}. \end{array} \tag{11}$$

For $a = 0$ symmetry X again has the form (10) where however $\lambda = \sigma_\mu = \omega_\mu = 0$. In addition, B^2 can depend not on t, x only, but also on u.

In accordance with (5), (7) and (9) equation (2) admits symmetry (10) iff the following classifying equations are satisfied:

$$\begin{array}{l} \big(\lambda(m+4)t + \mu + \frac{1}{a}\big(\frac{1}{2}\lambda x^2 + \sigma_\mu x_\mu + \gamma e^{\gamma t}\omega_\mu x_\mu\big)\big) + C^1\big) f^1 \\ +C^1_t u + B^1_t - a\Delta B^1 = (B^1\partial_u + B^2\partial_v + C^1(u\partial_u + v\partial_v) + C^2 u\partial_v \\ +\lambda m t(u\partial_u + v\partial_v) + \left(\frac{1}{2}\lambda x^2 + \sigma_\mu x_\mu + \gamma e^{\gamma t}\omega_\mu x_\mu\right)\left(\frac{1}{a}(u\partial_u + v\partial_v) - \frac{1}{a^2}u\partial_v\right)\big) f^1, \\ \\ (\lambda(m+4)t + \mu + C^1) f^2 + C^2 f^1 + \left(\frac{1}{2}\lambda x^2 + \sigma_\mu x_\mu + \gamma e^{\gamma t}\omega_\mu x_\mu\right)\left(\frac{1}{a} f^2 - f^1\right) \\ +C^1_t v + C^2_t u + B^2_t - \Delta B^2 - a\Delta B^1 = (B^1\partial_u + B^2\partial_v + C^1(u\partial_u + v\partial_v) + C^2 u\partial_v \\ +\lambda m t(u\partial_u + v\partial_v) + \left(\frac{1}{2}\lambda x^2 + \sigma_\mu x_\mu + \gamma e^{\gamma t}\omega_\mu x_\mu\right)\left(\frac{1}{a}(u\partial_u + v\partial_v) - \frac{1}{a^2}u\partial_v\right)\big) f^2. \end{array} \tag{12}$$

Equations (12) are nothing but a special case of the generic classifying equations (2.9), ref. [1] which are valid for arbitrary invertible diffusion matrix.

Equation (3) needs a particular analysis. For $p \neq 0$ the related symmetry operator (5) reduces to the form

$$X = X_0 + \tilde{X} \tag{13}$$

where

$$X_0 = \lambda\partial_t + \nu_\alpha \partial_{x_\alpha} + \Psi^{kl} x_k \partial_l,$$

$$\tilde{X} = \mu\left(3t\partial_t + 2x_\nu\partial_\nu - v\partial_v\right) - F\left(u\partial_u + v\partial_v\right) - B^1\partial_u - B^2\partial_v. \tag{14}$$

Here $\Psi^{\mu\nu}$ is an antisymmetric tensor and summation is imposed over the repeated Greek and Latin indices indices from 1 to m and from 1 to $m-1$ correspondingly.

Symmetries X_0 are admitted by equations (3), (4) with arbitrary non-linearities f^1 and f^2 while the classifying equations generated by symmetries $\tilde{X}$ are

$$\begin{aligned}
&(3\mu + F)f^1 + F_t u + B^1_t - pB^2_{x_m} \\
&= \left(B^1\partial_u + B^2\partial_v + Fu\partial_u + (F+\mu)v\partial_v\right) f^1, \\
&\\
&(4\mu + F)f^2 + F_t v + B^2_t - \Delta B^1 \\
&= \left(B^1\partial_u + B^2\partial_v + Fu\partial_u + (F+\mu)v\partial_v\right) f^2
\end{aligned} \tag{15}$$

where F and B^1, B^2 are unknown functions of t and t, x respectively.

The determining equations for symmetries of equation (3) with $p = 0$ are qualitatively different for the cases, when the number m of spatial variables $x_1, x_2, \cdots x_m$ is $m = 1$, $m = 2$ and $m > 2$. The related generator (5) has the form

$$\begin{aligned}
&X = \alpha D + \left(\textstyle\int (N-M)dt\right)\partial_t + 2mH^a\partial_{x_a} - \left(N + (m-2)H^a_{x_a}\right)u\partial_u \\
&- \left(M + (m+2)H^\alpha_{x_\alpha}\right)v\partial_v - B^1\partial_u - B^2\partial_v - B^3 u\partial_v
\end{aligned} \tag{16}$$

where summation from 1 to m is imposed over repeating indices, the Greek letters denote arbitrary parameters, M, N are functions of t, B^1, B^3 are functions of t, x, B^3 are functions of t, x, u, and

$$H^a = 2\lambda_b x_b x_a - x^2\lambda_a \text{ if } m > 2.$$

For $m = 2$ H^a are arbitrary functions satisfying the Caushy-Rieman conditions:

$$H^1_{x_1} = H^2_{x_2}, \quad H^1_{x_2} = -H^2_{x_1}.$$

In the case when $m = 1$ H^1 is a function of x and the sums with respect to a in (16) are reduced to the only terms.

The related classifying equations have the form

$$\begin{aligned}
&\left(\alpha + 2N - M + (m-2)H^a_{x_a}\right) f^1 + N_t u + B^1_t = (B^1\partial_u + B^2\partial_v \\
&+B^3 u\partial_v + \left(N + (m-2)H^a_{x_a}\right)u\partial_u + \left(M + (m+2)H^a_{x_a}\right)v\partial_v\big) f^1, \\
&\\
&\left(\alpha + N + (m+2)H^a_{x_a}\right) f^2 + B^3 f^1 + M_t v + B^3_t u + B^2_t - \Delta B_1 \\
&+(2-m)\Delta H^a_{x_a} u = \big(B^1\partial_u + B^2\partial_v + B^3 u\partial_v + \left(N + (m-2)H^a_{x_a}\right)u\partial_u \\
&+ \left(M + (m+2)H^a_{x_a}\right)v\partial_v\big) f^2.
\end{aligned} \tag{17}$$

We notice that in this case symmetry classification appears to be rather complicated and cumbersome. Nevertheless, the classifying equations can be effectively solved using the approach outlined in the following sections.

3 Classification of symmetries

Following [1] we specify basic, main and extended symmetries for the analyzed systems of reaction-diffusion equations.

Basic symmetries form the kernel of the main symmetry group and so are admitted by equation (2) with arbitrary non-linearities f^1 and f^2. They are generated by the following infinitesimal operators:

$$P_0 = \partial_t, \quad P_\lambda = \partial_{x_\lambda}, \quad J_{\mu\nu} = x_\mu \partial_{x_\nu} - x_\nu \partial_{x_\mu} \tag{18}$$

and correspond to shifts of independent variables and rotations of variables x_ν.

Main symmetries form an important subclass of general symmetries (10) which correspond to $\lambda = \sigma_\nu = \omega_\nu = 0$ and so have the following form

$$\tilde{X} = \mu D - C^1(u\partial_u + v\partial_v) - C^2 u\partial_v - B^1\partial_u - B^2\partial_v. \tag{19}$$

To describe all Lie symmetries admitted by equation (2) we follow the procedure outlined in [1] which includes the following steps:

- Finding all main symmetries (19), i.e., solving equations (12) for $\Psi^{\mu\nu} = \nu = \rho_\nu = \sigma_\nu = \omega_\nu = 0$:

$$\begin{aligned}
&(\mu + C^1)f^1 + C^1_t u_1 + B^1_t - a\Delta B^1 \\
&= \left(C^1(u\partial_u + v\partial_v) + C^2 u\partial_v + B^1\partial_u + B^2\partial_v\right) f^1, \\
&(\mu + C^1)f^2 + C^2_t u + C^1_t v + B^2_t - a\Delta B^2 - \Delta B^1 \\
&= \left(C^1(u\partial_u + v\partial_v) + C^2 u\partial_v + B^1\partial_u + B^2\partial_v\right) f^2.
\end{aligned} \tag{20}$$

- Specifying all cases when the main symmetries can be extended, i.e., at least one of the following systems is satisfied:

$$\begin{aligned}
&af^1 = \left(a(u\partial_u + v\partial_v) - u\partial_v\right) f^1, \\
&af^2 - f^1 = \left(a(u\partial_u + v\partial_v) - u\partial_v\right) f^2;
\end{aligned} \tag{21}$$

$$\begin{aligned}
&a(f^1 + \gamma u) = \left(a(u\partial_u + v\partial_v) - u\partial_v\right) f^1, \\
&a(f^2 + \gamma v) - \gamma u = \left(a(u\partial_u + v\partial_v) - u\partial_v\right) f^2
\end{aligned} \tag{22}$$

or if equation (21) is satisfied together with the following condition:

$$(m + 4)f^a = m(u\partial_u + v\partial_v)f^a, \quad a = 1, 2 \tag{23}$$

If relations (21), (22) or (23) are valid then the system (2) admits symmetry $G_\alpha, \widehat{G}_\alpha$ or K correspondingly.

- When classifying equations (3) or (2) with $a = 0$ the second step in not needed in as much as in accordance with (16) and (14) these equations admit basic and main symmetries only.

We note that the first step of the described procedure appears to be very complicated. In spite of that equations (20) can be effectively solved using separation of variables, the tree of versions of such separations is rather large and includes a lot of intersections.

In the next section we present specific tools used to overcome these difficulties.

4 Algebras of main symmetries for equation (2)

In accordance with the plane outlined in Section 4, to make symmetry classification of equations (2) we first describe the main symmetries generated by operators (19) and then indicate extensions of these symmetries.

First we note that for any f^1 and f^2 equation (2) admits the following equivalence transformations

$$\begin{array}{l} u \to K^1 u + b^1, \quad v \to K^1 v + K^2 u + b^2, \\ f^1 \to \lambda^2 K^1 f^1, \quad f^2 \to \lambda^2 (K^1 f^2 + K^2 f^1), \\ t \to \lambda^{-2} t, \quad x_b \to \lambda^{-1} x_b \end{array} \tag{24}$$

where K^1, K^2 and λ are constants which are distinct from zero, b^1 and b^2 are arbitrary constants. In accordance with its definition, equivalence transformations keep the general form of equation (2) but can change the concrete realization of their r.h.s.. For some non-linearities f^1 and f^2 there exist additional equivalence transformation which will be specified in the following.

We will use transformations (24) to simplify generators (19).

To solve rather complicated classifying equations (20), we use the main algebraic property of the main symmetries, i.e., the fact that they should form a Lie algebra (which we denote by $\mathcal{A}$). In other words, instead of going throw all non-equivalent possibilities arising via separation of variables in the classifying equations we first specify all non-equivalent realizations of algebra $\mathcal{A}$ for our equations up to arbitrary constants and arbitrary functions. Then we easily solve classifying equations (20) with *known functions* C^k *and* B^a.

Consider consequently one-, two-, $\cdots$ n-dimensional algebras of operators (25) which we write in the form

$$\tilde{X} = \mu D + N, \quad N = C^1 (u\partial_u + v\partial_v) + C^2 u\partial_v + B^1 \partial_u + B^2 \partial_v. \tag{25}$$

Let (25) be a basis element of a one-dimensional algebra $\mathcal{A}$ then commutators of $\tilde{X}$ with P_0 and P_a are equal to a linear combination of $\tilde{X}$ and operators (18). This can happen in the following cases:

$$\begin{array}{l} C^a = \mu^a, \ B^a = \nu^a; \quad C^a = e^{\lambda t} \mu^a, \ B^a = e^{\lambda t} \nu^a, \ \mu = 0; \\ C^a = 0, \ B^a = e^{\lambda t + \omega \cdot x} \nu^a, \ \mu = 0 \end{array} \tag{26}$$

where μ^{ab}, μ^a, λ, and $\omega = (\omega_1, \omega_2, \cdots, \omega_m)$ are constants, $\omega \cdot x = \omega_\nu x_\nu$ and $\omega \neq 0$.

To classify all non-equivalent symmetries (25), (26) we use isomorphism of the related N with 3×3 matrices of the following form

$$g = \begin{pmatrix} 0 & 0 & 0 \\ \nu^1 & \mu^1 & 0 \\ \nu^2 & \mu^2 & \mu^1 \end{pmatrix}. \tag{27}$$

Equivalence transformations (24) generate the following transformation for matrix (27)

$$g \to g' = UgU^{-1} \tag{28}$$

where

$$U = \begin{pmatrix} 1 & 0 & 0 \\ b^1 & K^1 & 0 \\ b^2 & K^2 & K^1 \end{pmatrix}, \qquad U^{-1} = \frac{1}{K^1} \begin{pmatrix} K^1 & 0 & 0 \\ -b^1 & 1 & 0 \\ b^1K^2 - b_2 & -\frac{K^2}{K^1} & 1 \end{pmatrix}. \tag{29}$$

Up to transformations (28) there exist six non-equivalent matrices g, i.e.,

$$\begin{gathered} g_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad g_2 = \begin{pmatrix} 0 & 0 & 0 \\ \lambda & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad g_3 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ g_4 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \quad g_5 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad g_6 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \end{gathered} \tag{30}$$

Let us denote

$$\hat{g} = g^{22}u\partial_u + g^{33}v\partial_v + g^{32}u\partial_v + g^{21}\partial_u + g^{31}\partial_v$$

where g^{ks} are elements of matrix g. Then in accordance with (26) the related symmetries (25) have the form

$$\begin{aligned} &\tilde{X} = \mu D + \hat{g} \ \text{ for } g = g_1, g_4, g_5, g_6, \\ &\tilde{X} = e^{\lambda t + \omega \cdot x}\tilde{g} \ \text{ for } g = g_1, g_2, \\ &\tilde{X} = e^{\lambda t}\hat{g} \text{ for any } g \text{ (30)}. \end{aligned} \tag{31}$$

Formulae (30)-(31) give the principal description of all possible one-dimension algebras $\mathcal{A}$ which can be admitted by equation (2).

To describe two-dimension algebras $\mathcal{A}$ we classify matrices g (27) forming two-dimension Lie algebras. Up to equivalence transformations (24) there exist six such algebras:

$$A_{2,1} = \{g_3, \tilde{g}_2\}, \ A_{2,2} = \{g_1, g_5\}, \ A_{2,3} = \{g_5, \tilde{g}_2\}, \ A_{2,4} = \{g_6, \tilde{g}_2\}, \tag{32}$$

$$A_{2,5} = \{g_1, g_2\},\ A_{2,13} = \{g_1, g_3\} \tag{33}$$

where $\tilde{g}_2$ is matrix g_2 (30) with $\lambda = 0$.

Algebras (32) are Abelian while (33) are characterized by the following commutation relations:

$$[e_1, e_2] = e_2. \tag{34}$$

Two-dimension algebras $\mathcal{A}$ generated by (32) and (33) are spanned on the following basis elements

$$\begin{array}{l} < \mu D + \hat{e}_1 + \nu t\hat{e}_2, \hat{e}_2 >,\ < \mu D + \hat{e}_2 + \nu t\hat{e}_1, \hat{e}_1 > \\ < \mu D - \hat{e}_1, \nu D - \hat{e}_2 >,\ < F_1\hat{e}_1 + G_1\hat{e}_2,\ F_2\hat{e}_1 + G_2\hat{e}_2 > \end{array} \tag{35}$$

and

$$< \mu D - \hat{e}_1, \hat{e}_2 >,\ < \mu D + \hat{e}_1 + \nu t\hat{e}_2, \hat{e}_2 > \tag{36}$$

respectively, where $\{F_1, G_1\}$ and $\{F_2, G_2\}$ are fundamental solutions of the following system

$$F_t = \lambda F + \alpha G, \quad G_t = \sigma F + \gamma G \tag{37}$$

with arbitrary parameters $\lambda, \alpha, \sigma, \gamma$. Arbitrary parameters μ and ν in particular can be equal to zero.

In addition, there exist two dimension algebras $\mathcal{A}$ which are induced by one-dimension algebras of matrices g (27), namely

$$< F\hat{g}, G\hat{g} >, \quad < \mu D + \lambda e^{\nu t + \omega\cdot x}\hat{g}, e^{\nu t+\omega\cdot x}\hat{g} > \tag{38}$$

with F and G satisfying (37). Such algebras correspond to incompatible classifying equations (20).

Up to transformations (28) there exist four three-dimension algebras $A_{3,1} - A_{3,4}$ of matrices (27) and the only four-dimension algebra of such matrices which we denote as A_4:

Table 1. Three- and four-dimension algebras of matrices (27)

Algebra	Basis elements	Nonzero commutators
$A_{3,1}$	$e_1 = g_1,\ e_2 = g_3,\ e_3 = \tilde{g}_2$	$[e_1, e_2] = e_2, \ \ [e_1, e_3] = e_3$
$A_{3,2}$	$e_1 = g_5,\ e_2 = g_1,\ e_3 = \tilde{g}_2$	$[e_2, e_3] = e_3$
$A_{3,3}$	$e_1 = \tilde{g}_2,\ e_2 = g_5,\ e_3 = g_6$	$[e_2, e_3] = e_1$
$A_{3,4}$	$e_1 = \tilde{g}_2,\ e_2 = g_3,\ e_3 = g_4$	$[e_1, e_2] = e_2,\ [e_1, e_3] = e_2 + e_3$
A_4	$e_1 = g_1,\ e_2 = g_3,\ e_3 = \tilde{g}_2,$ $e_4 = g_5$	$[e_1, e_2] = e_2,\ [e_1, e_3] = e_3,$ $[e_4, e_2] = e_3$

Using commutation relations present in the table we come to the following related three-dimension algebras $\mathcal{A}$:

$$< \mu D - \hat{e}_1,\ \hat{e}_2,\ \hat{e}_3 >,\ < \hat{e}_1,\ \ F_1\hat{e}_2 + G_1\hat{e}_3,\ F_2\hat{e}_2 + G_2\hat{e}_3 >$$

with e_a belonging to $A_{3,1}$;

$$< \mu D - 2\hat{e}_1,\ \nu D - 2\hat{e}_2, \hat{e}_3 >$$

with e_a belonging to $A_{3,2}$;

$$\begin{array}{l} < \mu D - 2\hat{e}_2,\ \nu D - 2\hat{e}_3,\ \hat{e}_1 >,\ < \hat{e}_1,\ D + 2e_\alpha + 2\nu t\hat{e}_1,\ \hat{e}_{\alpha'} >, \\ < e^{\nu t+\omega\cdot x}\hat{e}_1,\ e^{\nu t+\omega\cdot x}\hat{e}_\alpha,\ \hat{e}_{\alpha'} > \end{array}$$

where $\alpha,\ \alpha' = 2, 3, \alpha' \neq \alpha$ and e_a belong to $A_{3,3}$;

$$< \mu D - 2\hat{e}_1,\ \hat{e}_2,\ \hat{e}_3 >,\ < \hat{e}_1,\ e^{\nu t+\omega\cdot x}\hat{e}_2,\ \ e^{\nu t+\omega\cdot x}\hat{e}_3 >$$

with e_a belonging to $A_{3,4}$.

The four-dimensional algebra A_4 induces algebras $\mathcal{A}$ given below:

$$< \mu D - 2\hat{e}_1,\ \nu D - 2\hat{e}_4,\ \hat{e}_2,\ \hat{e}_3 >,\ \ < e^{\nu t+\omega\cdot x}\hat{e}_1,\ e^{\nu t+\omega\cdot x}\hat{e}_4,\ \hat{e}_2,\ \hat{e}_3 >\ .$$

Thus we had specified algebras of main symmetries which can be admitted by equation (2).

5 Solution of classifying equations for the case of invertible diffusion matrix

Applying results of the previous section we can easily classify main symmetries of equation (2). Such classification reduces to solving equations (20) with their *known coefficients* C^1, C^2 and B^1, B^2 which can be found comparing (19) with the found realizations of algebras $\mathcal{A}$. To complete the group classification of equations (2) we will specify all cases when relations (21)-(22) are satisfied, i.e., when the main symmetries can be extended.

Solving of equations (20) with known C^1, C^2 and B^1, B^2 is a rather routine procedure. We restrict ourselves to presentation of an example of such solution in the following. Note that asking for invariance of (2) w.r.t. one-dimension algebra $\mathcal{A}$ we fix the non-linearities f^1 and f^2 up to arbitrary functions while an invariance w.r.t. a two-dimension algebra $\mathcal{A}$ usually fixes these non-linearities up to arbitrary parameters.

We will solve classifying equations (20) up to equivalence transformations $U \to \tilde{U} = G(U, t, x)$, $t \to \tilde{t} = T(U, t, x)$, $x \to \tilde{x} = X(U, t, x)$ and $f \to \tilde{f} =$

$F(U, t, x, f)$ which keep the general form of equations (2) but can change functions f^1 and f^2.

The group of equivalence transformations for equation (2) can be found using the classical Lie approach and treating f^1 and f^2 as additional dependent variables. In accordance with their definition, equivalence transformations include all symmetry transformations (i.e., transformations generated by operators (18) and other symmetries which will be found in the following) and also transformations (24). In addition, for some particular non-linearities f^1 and f^2 there exist additional equivalence transformations, whose list is given in formulae (39):

$$\begin{array}{ll}
1. & u \rightarrow \exp(\omega t)u, \quad v \rightarrow \exp(\omega t)v, \\
2. & u \rightarrow u + \omega t + \mu x^2, \ v \rightarrow v, \\
3. & u \rightarrow u, \ v \rightarrow v + \rho t + \mu x^2, \\
4. & u \rightarrow u + \rho t, \ v \rightarrow v \exp(\rho t), \\
5. & u \rightarrow u, \ v \rightarrow v + \rho t u, \\
6. & u \rightarrow \exp(\omega t)u, \ v \rightarrow v + \kappa t u + \rho \frac{t^2}{2}, \\
7. & u \rightarrow u, \ v \rightarrow v - \rho t u + \rho \lambda \frac{t^2}{2}, \\
8. & u \rightarrow \exp(\rho t)u, \ v \rightarrow \exp(\rho t) \left(v + \varepsilon \frac{t^2}{2} u \right), \\
9. & u \rightarrow u + \rho t, \ v \rightarrow v + \rho t u + \rho \frac{t^2}{2}, \\
10. & u \rightarrow \exp(\omega t)u, \ v \rightarrow \exp(\omega t)(v - \omega t u), \\
11. & \texttt{Transformations (48) valid for } a = 0 \text{ only.}
\end{array} \tag{39}$$

where $\Phi(u)$ is an arbitrary function of u, F_1 and F_2 are functions of u which appear in the classified equations.

In the following we specify additional equivalence transformations (39) admitted by some of equations (2).

Let us present an example of solving of classifying equations. Consider the first of algebras $\mathcal{A}$ given by relation (35) with $e_1, \ e_2$ belonging to algebra $A_{2,2}$ (32). It includes two basis elements

$$X_1 = \mu D - u\partial_u - v\partial_v, \quad X_2 = \nu D - u\partial_v.$$

Comparing X_1 with $\tilde{X}$ (25) we conclude that in this case $C^1 = 1, C^2 = B^1 = B^2 = 0$ and so the classifying equations (20) are reduced to the following ones:

$$(\mu + 1) f^a = (u\partial_u + v\partial_v) f^a, \quad a = 1, 2$$

General solutions of this system have the form

$$f^1 = u^{\mu+1} F_1, \quad f^2 = v^{\mu+1} F_2 \tag{40}$$

where F_1 and F_2 are arbitrary functions of $\frac{u}{v}$.

Thus equation (2) admits symmetry X_1 iff the related non-linearities are of the form (40).

Asking for symmetry of equation (2) w.r.t. transformations generated by X_2 we come to the following classifying equations (20):

$$\nu f^1 = u\partial_v f^1, \quad \nu f^2 + f^1 = u\partial_v f^2. \tag{41}$$

Substituting (40) into (41) we come to the equation whose general solution for $\mu \neq 0$ is

$$f^1 = \lambda u^{\mu+1} e^{\nu\frac{v}{u}}, \quad f^2 = e^{\nu\frac{v}{u}}(\lambda v + \sigma u)u^{\mu}. \tag{42}$$

In the special case $\mu = 0$ the solution has the form

$$f^1 = \lambda u e^{\nu\frac{v}{u}} + \omega u, \quad f^2 = e^{\nu\frac{v}{u}}(\lambda v + \sigma u) + \omega v. \tag{43}$$

However, in this case there exist the additional equivalence transformation (39) given in Item 1 with $\rho = \omega$, which reduces parameter ω in (43) to zero. So without loss of generality we can restrict ourselves to solutions (42) for any μ.

We see that equation (2) admits the two-dimension algebra of main symmetries spanned on X_1, X_2 provided f^1 and f^2 have the form (42). This symmetry can be extended if functions (42) and f^2 satisfy one of conditions (21), (22) or both the conditions (21), (23).

Equation (22) is incompatible with (42). In order equation (21) be satisfied we have to impose the condition $\mu = -a\nu$ on parameters μ, ν and a. The related equation (2), (42) has the form

$$\begin{aligned} &u_t - a\Delta u = f^1 = \lambda u^{1-a\nu} e^{\nu\frac{v}{u}}, \\ &v_t - \Delta u - a\Delta v = e^{\nu\frac{v}{u}}(\lambda v + \sigma u)u^{-a\nu} \end{aligned} \tag{44}$$

and admits the Galilei generators G_μ (11). Finally, asking for equation (23) be satisfied we obtain one more condition $\nu = -\frac{4}{a}m$ which guaranties invariance of equation (44) w.r.t. the conformal generator K of (11).

The obtained classification results are presented in Table 2, Item 3.

In analogous way we solve classifying equations for other algebras $\mathcal{A}$ and specify the cases when the main symmetry can be extended.

6 Classification results for equations (2) with invertible diffusion matrix

The classification results are given in the following Tables 1-5 when we also indicate the additional equivalence transformations (AET) (39) which are admitted by some particular equations (2). The symbols D, $\hat{G}_\nu$, G_ν and K are used to denote operators (11). In addition, we denote

$$\tilde{K} = K + \frac{1}{\lambda - 1}\left(t\left(pu\partial_u + (2-\lambda)v\partial_v\right) + u\partial_v\right).$$

To save a room we present in Table 2, Items 8-11 and Table 3, Items 1-3, the classification results which are valid for equation (2) with $a \neq 0$ and $a = 0$ as well. The completed analysis of symmetries and the corresponding non-linearities for equation (2) with $a = 0$ is given in Section 7.

In the following tables F_1, F_2 and F are arbitrary functions whose arguments are specified in the third column, $\Psi(x)$ is an arbitrary function of $x_1,\ x_2,\ \cdots,\ x_m$ and $\psi_\nu(x)$ is a solution of the linear heat equation $(\partial_t - \Delta)\psi_\nu = \nu\psi_\nu$. In addition, we denote by $\Psi_\mu(x)$ a solution of the Laplace equation $\Delta\Psi_\mu(x) = \mu\Psi_\mu(x)$.

Greek letters in the tables denote arbitrary parameters which can take any (including zero) real values. The only exception is parameter ε in as much as without loss of generality we can restrict ourselves to its values $\varepsilon = \pm 1$.

Table 2. Non-linearities with arbitrary functions and symmetries for equations (2)

No	Nonlinear terms	Arguments of F_1, F_2	Symmetries
1.	$f^1 = \nu u + F_1$, $f_2 = \nu u^2 + F_1 u + F_2$,	$2v - u^2$	$\psi_\nu\,(u\partial_v + \partial_u)$
2.	$f^1 = e^{\nu u} F_1$, $f^2 = e^{\nu u}(F_2 + F_1 u)$,	$2v - u^2$	$\nu D - u\partial_v - \partial_u$
3.	$f^1 = F_1$, $f^2 = F_2 + \nu v$,	u	$\psi_\nu \partial_v$
4.	$f^1 = \alpha u + \mu$, $f_2 = \nu v + F,\ \alpha\mu = 0$	u	$\psi_\nu\partial_v$, $e^{(\nu-\alpha)t}\,(u - \mu t)\,\partial_v$
5.	$f^1 = u^2$, $f^2 = uv + \nu v + F$	u	$e^{\nu t} u\partial_v$, $e^{\nu t}\,(\partial_v + tu\partial_v)$
6.	$f^1 = (u^2 - 1)$, $f^2 = (u + \nu)\,v + F$	u	$e^{(\nu+1)t}\,(u\partial_v + \partial_v)$, $e^{(\nu-1)t}\,(u\partial_v - \partial_v)$
7.	$f^1 = (u^2 + 1)$, $f_2 = (u + \nu)\,v + F$	u	$e^{\nu t}\,(\cos tu\partial_v - \sin t\partial_v)$, $e^{\nu t}\,(\sin tu\partial_v + \cos t\partial_v)$
8*.	$f^1 = e^{\nu v} F_1,\ f^2 = e^{\nu v} F_2$	u	$\nu D - \partial_v$
9*.	$f^1 = e^{\nu u} F_1$, $f^2 = e^{\nu u} F_2$	v	$\nu D - \partial_u$
10*.	$f^1 = \nu u + F_1$, $f_2 = -a\mu u + F_2$	v	$e^{(\nu + a\mu)t}\Psi_\mu(x)\partial_u$
11*.	$f^1 = u(F_1 + \nu \ln u)$, $f^2 = v(F_2 + \nu \ln u),\ \nu \neq 0$	$\frac{u}{v}$	$e^{\nu t}(u\partial_u + v\partial_v)$

The items marked by asterisks are valid for both cases $a \neq 0$ and $a = 0$. If $a = 0$ then in Item 8* without loss of generality $F_1 = 1$.

Table 3. Non-linearities with arbitrary functions, symmetries and AET for equations (2)

No	Nonlinear terms	Arguments of F_1, F_2	Symmetries	AET (39)
1*.	$f^1 = uF_1 - \nu v,$ $f^2 = \nu\frac{v}{u}(v-u)$ $+uF_2 - vF_1,\ \nu \neq 0$	$ue^{\frac{v}{u}}$	$e^{\nu t}\,(u\partial_v - u\partial_u$ $-v\partial_v)$ & $\widehat{G}_\alpha$ if $a = 1$	
2*.	$f^1 = u^{\nu+1}F_1,$ $f^2 = u^\nu\,(F_2 u - F_1 v)$	$ue^{\frac{v}{u}}$	$\nu D + u\partial_v - u\partial_u$ $-v\partial_v$ & G_α if $\nu = 0,\ a = 1$	
3*.	$f^1 = u^{\mu+1}F_1,$ $f^2 = u^{\mu+1}F_2$	$\frac{v}{u}$	$\mu D - u\partial_u - v\partial_v$	$1,$ if $\mu = 0$
4.	$f^1 = e^{\nu\frac{v}{u}}F_1 u,$ $f^2 = e^{\nu\frac{v}{u}}\,(F_1 v + F_2)$	u	$\nu D - u\partial_v$	5 if $\nu = 0$
5.	$f^1 = u(F_1 - \nu),$ $f^2 = F_1 v + F_2,$ $\nu \neq 0$	u	$e^{\nu t}u\partial_v$ & $\psi_\nu\partial_v$ if $F_1 = \nu$	$3,$ if $F_1 = 0$

Table 4. Non-linearities with arbitrary parameters and extendible symmetries for equations (2) with $a \neq 0$

No	Nonlinear terms	Main symmetries	Additional symmetries	AET (39)
1.	$f^1 = \lambda u,$ $f^2 = \sigma u^\mu$	$\psi_0\partial_v,$ $e^{-\lambda t}u\partial_v$	$e^{\lambda t}\,(u\partial_v + \lambda\partial_u)$ if $\mu = 2$	3; 5 if $\lambda = 0$
2.	$f^1 = \lambda e^u,$ $f^2 = \sigma e^u$	$D - \partial_u,$ $\psi_0\partial_v$	$u\partial_v$ if $\lambda = 0$	3; 5 if $\lambda = 0$
3.	$f^1 = \lambda u^{\nu+1}e^{\mu\frac{v}{u}},$ $f^2 = e^{\mu\frac{v}{u}}(\lambda v + \sigma u)u^\nu$	$\mu D - u\partial_v,$ $\nu D - u\partial_u$ $-v\partial_v$	G_α if $\nu = a\mu$ & K if $\nu = \frac{4}{m}$	$1,$ if $\nu = 0;$ $5,$ if $\mu = 0$
4.	$f^1 = \lambda u^{\mu+1},$ $f^2 = \sigma u^{\mu+1},$ $\lambda\sigma = 0$	$\mu D - u\partial_u$ $-v\partial_v,$ $\psi_0\partial_v$	$u\partial_v$ if $\lambda = 0$	3; 5 if $\lambda = 0$

Table 5. Non-linearities with arbitrary parameters and non-extendible symmetries for equations (2) with $a \neq 0$

	Nonlinear terms	Symmetries	AET (39)
1.	$f^1 = \lambda v^{\nu+1}$, $f^2 = \mu v^{\nu+1}$	$\nu D - u\partial_u - v\partial_v$, $\Psi_0(x)\partial_u$	2
2.	$f^1 = \lambda e^v,\ f^2 = \sigma e^v$	$D - \partial_v,\ \Psi_0(x)\partial_u$	2
3.	$f^1 = \lambda e^u,\ f^2 = \sigma u e^u$	$D - \partial_u - 2u\partial_v,\ \psi_0\partial_v$	3; 5 if $\lambda = 0$
4.	$f^1 = \nu e^{\lambda\left(2v-u^2\right)}$, $f^2 = (\nu u + \mu)\, e^{\lambda\left(2v-u^2\right)}$	$\lambda D - \partial_v,\ \partial_u + u\partial_v$	9
5.	$f^1 = \mu \ln v$, $f^2 = \nu \ln v$	$\Psi_0(x)\partial_u,\ D + u\partial_u + v\partial_v$ $+ \left((\mu - \nu a)t - \frac{\nu}{2m} m x^2\right)\partial_u$	2
6.	$f^1 = \lambda$ $f^2 = \varepsilon \ln u$,	$D + u\partial_u + v\partial_v + \varepsilon t\partial_v$, $\psi_0\partial_v,\ (u - \lambda t)\,\partial_v$	3, 7; & 5 `if` $\lambda = 0$
7.	$f^1 = \lambda u^{\nu+1}$, $f^2 = \lambda u^{\nu+1}\ln u$, $\nu \neq -1$	$\nu D - u\partial_u - v\partial_v - u\partial_v$, $\psi_0\partial_v$	3
8.	$f^1 = \varepsilon(2v - u^2)$, $f^2 = (\mu + \varepsilon u)\,(2v - u^2)$ $-\varepsilon\frac{\mu^2}{2}u,\quad \mu \neq 0$	$X_1 = e^{\mu t}\,(2\partial_u + 2u\partial_v$ $+\mu\varepsilon\partial_v)\,,\quad 2tX_1 + \varepsilon e^{\mu t}\partial_v$	
9.	$f^1 = \varepsilon(2v - u^2)$, $f^2 = (\mu + \varepsilon u)\,(\varepsilon v - u^2)$ $+\varepsilon\frac{1-\mu^2}{2}u$	$X^{\pm} = e^{\mu\pm 1}\,(2\partial_u + 2u\partial_v$ $+\varepsilon(\mu \pm 1)\partial_v)$	9, if $\mu^2 = 1$
10.	$f^1 = \varepsilon(2v - u^2)$, $f^2 = (\mu + \varepsilon u)\,(2v - u^2)$ $-\varepsilon\frac{1+\mu^2}{2}u$	$e^{\mu t}\,(2\cos t\,(\partial_u + u\partial_v)$ $+\varepsilon(2\mu\cos t - \sin t)\partial_v)\,,$ $e^{\mu t}\,(2\sin t\,(\partial_u + u\partial_v)$ $+\varepsilon(2\mu\sin t + \cos t)\partial_v)$	

In the following table we present symmetries of a special subclass of equations (2) whose r.h.s. is given in the table title. The related non-linearities had appeared in the classification procedure of reaction-diffusion equations with the unit diffusion matrix [2] where they corresponded to a very rich spectrum of symmetries.

In the case of triangular diffusion matrices studied in the present paper the classification results present in Table 5 include thirteen non-equivalent types of equations. Among them there are five equations invariant w.r.t. two dimensional algebras of main symmetries, and four equations admitting symmetries G_ν i.e., being invariant w.r.t. the Galilei group.

In Table 5 the following notations are used: $\delta = \frac{1}{4}(\mu - \nu)^2 + \lambda\sigma$, $\omega_0 = \frac{1}{2}(\mu + \nu)$, $\omega_\pm = \omega_0 \pm 1$ where μ, ν, λ and σ are parameter used in definition of the related non-linearities f^1, f^2.

Table 6. Symmetries of equations (2) with non-linearities
$f^1 = \lambda v + \mu u \ln u$, $f^2 = \lambda \frac{v^2}{u} + (\sigma u + \mu v)\ln u + \nu v$ and $a \neq 0$

No	Conditions for coefficients	Main symmetries	Additional symmetries
1	$\lambda = 0$, $\mu \neq \nu$	$e^{\nu t} u \partial_v$,	$\psi_\nu \partial_v$ if $\mu = 0$, & $G_a \texttt{if}\ \nu = a\sigma$
		$e^{\mu t}\left((\mu - \nu) R\partial_R + \sigma u \partial_v\right)$	$\hat{G}_a$, if $\mu \neq 0$, $a\sigma = \nu - \mu$,
2	$\sigma = 0$, $\mu\lambda \neq 0$, $\mu \neq \nu$	$e^{\nu t}\left(\lambda R\partial_R + (\mu - \nu) u \partial_v\right)$	G_a if $\nu = 0$, $a\mu = -\lambda$
		$e^{\mu t} R\partial_R$	$\hat{G}_a$, if $a(\nu - \mu) = \lambda$
3	$\delta = 0$	$X_4 = e^{\omega_0 t}\left(2\lambda R\partial_R\right.$ $\left. + (\nu - \mu) u\partial_v\right)$,	G_a, if $\omega_0 = 0$, $a\nu = -\lambda \neq 0$
		$2e^{\omega_0 t} u \partial_v + tX_4$	$\widehat{G}_a$, if $\omega_0 \neq 0$, $2\lambda = a(\mu - \nu) \neq 0$
4	$\lambda \neq 0$, $\delta = 1$	$e^{\omega_+ t}\left(\lambda R\partial_R + (\omega_+ - \mu) u\partial_v\right)$,	G_a, if $a\mu = \lambda$ $\omega_+\omega_- = 0$
		$e^{\omega_- t}\left(\lambda R\partial_R + (\omega_- - \mu) u\partial_v\right)$	$\hat{G}_\alpha$, if $\omega_+ \neq 0$, $\lambda = -a(\omega_+ - \mu)$
5	$\lambda \neq 0$, $\delta = -1$	$e^{\omega_0 t}[2\lambda \cos t R\partial_R$ $+((\nu - \mu)\cos t - 2\sin t) u\partial_v]$, $e^{\omega_0 t}[2\lambda \sin t R\partial_R$ $+((\nu - \mu)\sin t + 2\cos t) u\partial_v]$	none

If $\lambda = \mu = 0$ or $\lambda = \nu = 0$ then the related equation (2) admits additional equivalence transformations 10 or 5 from the list (39) correspondingly.

We see that there exist a number of non-equivalent systems (2) with non-degenerate diffusion matrix. Rather surprisingly the number of such equation with nilpotent diffusion matrix (which are classified in the following section) appears to be even more large.

7 Group classification of reaction-diffusion equations with nilpotent diffusion matrix

7.1 Equations with first derivatives in x

Consider now equations (3), (4) and specify their Lie symmetries. In this subsection we restrict ourselves to the case $p \neq 0$ when generators of admitted Lie group have the general form (13) while the related classifying equations are given by formula (15). Moreover without loss of generality we put $p = 1$.

We solve the classifying equations using the technique developed in Sections 5 and 6. The general analysis of admissible algebras $\mathcal{A}$ can be carried out in complete analogy with Section 4. Moreover, the results present in Section 4 can be extended to the case of equations (3), (4) provided we make a formal change $D \to \tilde{D} = 3t\partial_t + 2x_\nu\partial_\nu - v\partial_v$ in all formulae where the operator D appeared, and exclude all algebras $\mathcal{A}$ where matrices g_4, g_5 and g_6 (30) appear. Of course it is necessary to take into account that in contrast with D operator $\tilde{D}$ does not commute with ∂_v. As a result we come to the following one-dimension algebras

$$\begin{aligned}
&\tilde{X}_1^{(1)} = \mu\tilde{D} - u\partial_u - v\partial_v,\\
&\tilde{X}_1^{(2)} = \tilde{D} - \nu\partial_u, \quad \tilde{X}_2^{(\nu)} = e^{\nu t}\left(u\partial_u + v\partial_v\right),\\
&\tilde{X}_1^{(3)} = \tilde{D} + u\partial_u + v\partial_v + \nu\partial_v, \quad \tilde{X}_3^{(3)} = e^{\sigma_3 t + \rho_3 x}\left(\partial_u + \partial_v\right),\\
&\tilde{X}_3^{(1)} = e^{\sigma_1 t + \rho_1\cdot x}\partial_u, \quad \tilde{X}_3^{(2)} = e^{\sigma_2 t + \rho_2\cdot x}\partial_v
\end{aligned} \tag{45}$$

and two-dimension algebras

$$\begin{aligned}
&\tilde{A}_1 =< \tilde{D}, \tilde{X}_2^{(0)} >, \ \tilde{A}_2 =< \tilde{X}_1^{(2)}, X_3^{(3)} >, \ \tilde{A}_3 =< \tilde{X}_1^{(3)}, \tilde{X}_3^{(1)} >,\\
&\tilde{A}_4 =< \tilde{X}_1^{(1)}, \tilde{X}_3^{(1)} >, \ \tilde{A}_6 =< \tilde{D} + 4(u\partial_u + v\partial_v) + t\partial_v, \quad X_3^{(2)} >,\\
&\tilde{A}_5 =< \tilde{X}_1^{(1)}, \tilde{X}_3^{(2)} >, \ \tilde{A}_7 =< \tilde{D} + 3(u\partial_u + v\partial_v) + t\partial_u, \quad X_3^{(1)} > .
\end{aligned} \tag{46}$$

In this way the problem of group classification of the equations with the first order derivative terms reduces to solving the classifying equations (15) which their known coefficients B^1, B^2, F and specifying the case when these equations have non-trivial solutions. These coefficients are easily identified comparing (45), (46) with (14). For example, for symmetry $\tilde{X}_1^{(1)}$ we have $F = 1, B^1 = B^2 = 0$, for $\tilde{X}_1^{(2)}$ the values of these coefficients are $B^1 = \nu, B^2 = F = 0, \mu = 1$, etc. Solving the related classifying equations (15) we easily find the related non-linearities f^1, f^2 which are given in Table 7.

In accordance with the results present in Table 7 equations (3), (4) with $p \neq 0$ can admit neither Galilei nor conformal symmetry transformations. This result follows directly from formulae (13), (14). equations.

In six cases enumerated in the table the corresponding equations (3), (4) admit infinite dimension symmetry algebras whose generators are defined up to arbitrary functions or arbitrary solutions of linear equations, see Items 5-7, 9-14 here. However, these infinite symmetries generate the same classifying equations as one dimensional algebras $X_3^1 - X_3^{(3)}$ and two dimensional algebras $\tilde{A}_3, \tilde{A}_4, \tilde{A}_7$.

Finally we note that in addition to their symmetries, equations (3), (4) with $p \neq 0$ admit equivalence transformations (24) with $K^2 = 0$. These transformations which change functions f^1, f^2 and also the value of parameter p where used to simplify generators (45), (46) and the corresponding equations (3), (4). Equations (3), (4) with the non-linearities given in Items 3 (when $\nu = 0$) and 8 of Table 6 admit additional equivalence transformation $u \to e^{\sigma t}u, \quad v \to e^{\sigma t}v$.

Table 7. Non-linearities and symmetries for equations (3), (4) with $p = 1$

	Non-linearities	Arguments of F_1 F_2	Symmetries
1.	$f^1 = u_1^{1+3\mu} F_1,$ $f^2 = u^{1+4\mu} F_2$	$vu^{-\mu-1}$	$2\mu\tilde{D} - u\partial_u - v\partial_v$
2.	$f^1 = v^3 F_1,$ $f^2 = u_2^4 F_2$	$u - \nu \ln v$	$\tilde{D} - \nu\partial_u$
3.	$f^1 = u(F_1 + \nu \ln u),$ $f^2 = v(F_2 + \nu \ln u)$	$\frac{v}{u}$	$e^{\nu t}\left(u\partial_u + v\partial_v\right)$
4.	$f^1 = u_1^{-2} F_1,$ $f^2 = u_1^{-3} F_2$	$v - \nu \ln u$	$\tilde{D} + u\partial_u + v\partial_v + \nu\partial_v$
5.	$f^1 = \lambda u + F_1,$ $f^2 = -\mu u + F_2$	v	$e^{\lambda t}\Psi_\mu(x)\partial_u$
6.	$f^1 = \nu v + F_1,$ $f^2 = \lambda v + F_2$	u	$e^{\lambda t - \nu x_m}\Psi(\tilde{x})\partial_v$
7.	$f^1 = \lambda u + F_1$ $f^2 = \sigma v + F_2$	$u - v$	$e^{\lambda t}\ e^{\frac{x_m+t}{2}}\Psi_\mu(\tilde{x}, x_m + t)\left(\partial_u + \partial_v\right),$ $\mu = \lambda - \sigma + \frac{1}{4}$
8.	$f^1 = \alpha u_1^{-2} u_2^3,$ $f^2 = \nu u_1^{-3} u_2^4$		$\tilde{D}, \quad u\partial_u + v\partial_v$
9.	$f^1 = \alpha e^{3u},$ $f^2 = \nu e^{4u}$		$\tilde{D} - \partial_u,\ \Psi(\tilde{x})\partial_v$
10.	$f^1 = \alpha e^{-2v},$ $f^2 = \nu e^{-3v}$		$\tilde{D} + u\partial_u + v\partial_v + \partial_v,\ \Psi_0(x)\partial_u$
11.	$f^1 = \alpha u^{3\mu+1},$ $f^2 = \nu u^{4\mu+1}$		$\mu\tilde{D} - u\partial_u - v\partial_v, \quad \Psi(\tilde{x})\partial_v$
12.	$f^1 = \alpha v^{2\nu+1},$ $f^2 = \nu v^{3\nu+1}$		$\nu\tilde{D} - u\partial_u - v\partial_v,\ \Psi_0(x)\partial_u$
13.	$f^1 = \alpha u^{\frac{1}{4}},$ $f^2 = \nu \ln u$		$\frac{1}{4}\tilde{D} + v\partial_v + u\partial_u + \nu t\partial_v,\ \Psi(\tilde{x})\partial_v$
14.	$f^1 = \nu \ln v,$ $f^2 = \frac{\alpha}{\sqrt{v}}$		$\frac{1}{3}\tilde{D} + u\partial_u + v\partial_v + \nu t\partial_u,\ \Psi_0(x)\partial_u$

Here $\Psi_\mu(x)$ and $\Psi(\tilde{x}, x_m + t)$ are arbitrary solutions of the Laplace equation $\Delta\Psi_\mu = \mu\Psi_\mu$ in m-dimensional space, $\tilde{\Psi}_\mu(\tilde{x})$ is a solution of the Laplace equation in $m-1$-dimensional space, $\tilde{x} = (x_1, x_2, \cdots, x_{m-1})$, μ, ν and λ are arbitrary parameters satisfying $\nu\lambda \neq 0$. Finally, we denote $\tilde{D} = 3t\partial_t + 2x_\nu\partial_\nu - v\partial_v$.

7.2 Equations (2) with $a = 0$

The procedure of classification of equations (3), (4) with $p = 0$ (or equations (2) with $a = 0$) appears to be more complicated then in the case of p non-zero. The general form of symmetry admitted by this equation is given by equation (16) while the classifying equations take the form (17).

A specific property of symmetries (16) is that in contrast with operators (10) and (13) they can generate non-linear transformations for dependent variables since the coefficient B^3 can be a function of u. In addition, the classifying equations critically depend on the number m of independent variables x_ν and are qualitatively different for the cases $m = 1, m = 2$ and $m > 2$. Nevertheless, these equations can be effectively solved with using the analysis of low dimension symmetry algebras present in Section 4.

One more specific point in the classification of equations (2) with $a = 0$ is that they admit powerful equivalence relations

$$u \to u, \ v \to v + \Phi(u) \tag{47}$$

and

$$u \to u, \ v \to v + \hat{\Phi}(u, t, x) \tag{48}$$

which did not appear in our analysis presented in the previous sections.

Equivalence transformation (47) (with $\Phi(u)$ being an arbitrary function of u) are admitted by any equation (2) with $a = 0$. Transformations (48) are valid for the cases when f^1 does not depend on v and in the same time f^2 is either linear in v or does not depend on this variable. Moreover, the related functions $\hat{\Phi}(u, t, x)$ should satisfy the following system of equations

$$\begin{array}{l} f^2_v\hat{\Phi}_t - \hat{\Phi}_{tt} - f^1\hat{\Phi}_{tu} = 0, \\ f^2_v\hat{\Phi}_{x_\nu} - \hat{\Phi}_{tx_\nu} - f^1\hat{\Phi}_{ux_\nu} = 0 \end{array} \tag{49}$$

Thus the group classification of equation (2) with $a = 0$ is reduced to solving the classifying equations (17) with using the algorithm presented in Section 3. We will not reproduce here cumbersome and rather routine calculations which are needed to classify equations (2) with $a = 0$ which can be carried out in the same way as in Sections 2-6 and taking into account the specific points mentioned in the above. The classification results are presented below in Tables 8-10 and also in Table 2, Items 8-11 and Table 3, Items 1-3. The related items are marked by asterisks. The additional equivalence transformations are specified below the Tables 8,9 and 10.

In Tables 8-10 the symbol W denotes a function of t, x and u which solve the following equation:

$$f^2_v - W_t - W_u f^1 = 0.$$

Table 8. Non-linearities with arbitrary functions for equations (2) with nilpotent diffusion matrix

No	Nonlinear terms	Arguments of F_α	Symmetries
1.	$f^1 = F_1 u^{\mu-\nu},\ f^2 = F_2 u^\mu$	$\frac{u^{\nu+1}}{v}$	$Q_1 = (\mu-1)D - \nu t\partial_t - u\partial_u$ $-(\nu+1)v\partial_v$ & $(m-2)x^2\partial_{x_a}$ $-x_a Q_1$ `if` $\nu(m-2) = 4$, $\mu(m-2) = m+2,\ m \neq 2$
2.	$f^1 = F_1 u v^{\mu-1},\ f^2 = F_2 v^\mu,$ $F_2 \neq 0$	u	$\mu D - t\partial_t - v\partial_v$ & $e^W\partial_v$ `if` $\mu = 1$ & $H^a\partial_{x_a} - H^b_{x_b} v\partial_v$ `if` $m = 2$
3.	$f^1 = F_1 v^{-1},\ f^2 = F_2 + \nu v$	u	$e^{\nu t}\left(\partial_t + \nu v\partial_v\right)$ & $e^W\partial_v$ `if` $F_1 = 0$
4.	$f^1 = F_1 v^{\mu-1},\ f^2 = F_2 v^\mu$	ve^u	$\mu D - t\partial_t - v\partial_v + \partial_u$
5.	$f^1 = \frac{F_1}{v} + \nu,\ f^2 = F_2 + \nu v$	ve^u	$e^{\nu t}\left(\partial_t + \nu v\partial_v - \nu\partial_u\right)$
6.	$f^1 = 0,\ f^2 = F_2$	v	$\Psi_0(x)\partial_u,\ x_a\partial_{x_a} + 2u\partial_u$
7.	$f^1 = F_1,\ f^2 = 0$	u	$e^W\partial_v,\ \ x_a\partial_{x_a} - 2v\partial_v$
8.	$f^1 = \frac{\nu}{\mu-1}u + F_1 u^{2-\mu},$ $f^2 = \frac{\mu\nu}{\mu-1}v + F_2 u,\ \mu \neq 1$	$vu^{-\mu}$	$e^{\nu t}\left((1-\mu)t\partial_t - \nu u\partial_u\right.$ $\left.-\nu\mu v\partial_v\right)$
9.	$f^1 = uF_1,\ m = 1$ $f^2 = vF_2 + u$	vu^3	$Q_2 = \cos(2x)\left(u\partial_u - 3v\partial_v\right)$ $+\sin(2x)x\partial_x,\ \ Q_3 = (Q_2)_x$
10.	$f^1 = uF_1,\ m = 1$ $f^2 = vF_2 - u$	vu^3	$Q_4 = e^{2x}\left(\partial_x + u\partial_u - 3v\partial_v\right),$ $Q_5 = e^{-2x}\left(\partial_x - u\partial_u + 3v\partial_v\right)$
11.	$f^1 = F_1, f^2 = vF_2,\ m = 2$	ve^u	$H^a\partial_{x_a} - H^b_{x_b}\left(v\partial_v - \partial_u\right)$
12.	$f^1 = \nu e^{\frac{v}{u}},\ f^2 = e^{\frac{v}{u}}F$	u	$D - u\partial_v$
13.	$f^1 = F_1,\ F_2 = vF_2 + F_3$	u	$e^W\partial_v$

For the non-linearities enumerated in Items 2 (when $\mu = 1$), 3 (when $F_1 = 0$), 4 and 8 of Table 8 the related equation (2) with $a = 0$ admits additional equivalence transformations (48). In addition, transformations (24) and (47) and some equivalence transformations from the list (39) are admissible, namely, transformations (2) for the non-linearities given in Item 1 (when $\nu = -1, \mu = 0$) and in Item 6, transformations (1) and (3) for the non-linearities from Item I (when $\nu = 1, \mu = 0$) and Item 7 respectively.

Table 9. Non-linearities with arbitrary parameters and extendible symmetries for equations (2) with nilpotent diffusion matrix

No	Non-linearities	Main symmetries	Additional symmetries	AET (39)
1.	$f^1 = \lambda u^{\nu+1} v^{\mu}$, $f^2 = \sigma u^{\nu} v^{\mu+1}$	$(\mu+\nu)t\partial_t$ $-(\mu+1)u\partial_u$ $+(\nu-1)v\partial_v$ $Q_6 = 2\mu u\partial_u$ $+(\mu+\nu)x_a\partial_{x_a}$ $-2\nu v\partial_v$	$x_a Q_6 - 2\kappa x^2\partial_{x_a}$ if $\kappa(m+2) = \nu$, $\kappa(2-m) = \mu$	
			$e^W\partial_v$ if $\lambda = 0$, $\mu = -1$ & $2x^2\partial_{x_a}$ $-(m-2)x_a Q_6$ if $\nu = \frac{m+2}{m-2}$, $m \neq 2$	11, $3, 5$
			$\Psi_0(x)\partial_u$, if $\sigma = 0, \nu = -1$, & $x_a Q_6 + \frac{2}{m+2} x^2\partial_{x_a}$ if $\mu = \frac{m-2}{m+2}$, $m \neq 2$	11, 2
			$\Psi_0(x)\partial_u$ if $\lambda = \nu = 0$	11, 2
			$e^W\partial_v$ if $\mu = 0$ & $H^a\partial_{x_a} - H^a_{x_a} v\partial_v$ if $m = 2$	11 & 5 if $\lambda = \sigma$
2.	$f^1 = \lambda u^{\nu+1} v^{-1}$, $f^2 = \sigma u^{\nu} + \varepsilon v$, $\lambda \neq 0$	$e^{\varepsilon t}(\partial_t + \varepsilon v\partial_v)$, ${Q'}_6 = Q_6\|_{\mu=-1}$	$x_a {Q'}_6 - \frac{2}{m-2} x^2\partial_{x_a}$ if $\nu = \frac{m+2}{m-2}$, $m \neq 2$	
			$\Psi_0(x)\partial_u$ if $\sigma = 0$, $\nu = -1$	2
			$v\partial_v + u\partial_u$ if $\sigma = 0$, $\nu = 1$	1
3.	$f^1 = \lambda e^{\nu u}$, $f^2 = \sigma e^{(\nu+1)u}$, $\lambda\sigma = 0$	$(\nu+1)D - v\partial_v$ $-t\partial_t - \partial_u$, $e^W\partial_v$	$\nu(v\partial_v + t\partial_t) - \partial_u$ if $\sigma = 0$	11, 3
			$v\partial_v + t\partial_t$ if $\lambda = 0$	11, $3, 5$
4.	$f^1 = \lambda e^{(\nu+1)v}$, $f^2 = \sigma e^{\nu v}$	$(\nu-1)D - u\partial_u$ $+t\partial_t - \partial_v$, $\Psi_0(x)\partial_u$	$\nu(u\partial_u - t\partial_t)$ $+\partial_v$ if $\lambda = 0$	2 for any λ
5.	$f^1 = \lambda v^{\mu-1} e^u$, $f^2 = \sigma v^{\mu} e^u$	$D - \partial_u$, $t\partial_t + v\partial_v - \mu\partial_u$	e^W if $\mu = 1$ & $H^a\partial_{x_a} - H^a_{x_a} v\partial_v$ if $m = 2$	11 if $\mu = 1$
6.	$f^1 = \lambda \ln v$, $f^2 = \sigma v^{\frac{\mu+1}{2}}$	$\mu D - \frac{\mu+1}{2} t\partial_t$ $-\frac{1-\mu}{2} u\partial_u - v\partial_v$ $-\lambda t\partial_u$, $\Psi_0(x)\partial_u$	$x_a\partial_{x_a} + 2u\partial_u$, $u\partial_u + 2v\partial_v + t\partial_t$ $+2\lambda t\partial_u$ if $\sigma = 0$	2
7.	$f^1 = \sigma v^{1-\mu}$, $f^2 = \lambda \ln v$	$\mu D + \mu u\partial_u - v\partial_v$ $-(1+\mu)t\partial_t$ $-\frac{\lambda x^2}{2m}\partial_u$, $\Psi(x)\partial_u$	$D + u\partial_u - t\partial_t$ if $\sigma = 0$	2

Table 10. Non-linearities with arbitrary parameters and non extendible symmetries for equations (2) with $a = 0$

No	Non-linearities	Conditions	Symmetries	AET (39)
1.	$f^1 = \lambda u^{3\mu+1} v^{\mu}$, $f^2 = \sigma u^{3\mu} v^{\mu+1} - \alpha u$,	$\mu \neq 0$, $m = 1$, $\alpha = -1$	$Q_7 = 4\mu t\partial_t$ $-(\mu+1)u\partial_u$ $+(3\mu-1)v\partial_v$, Q_2, Q_3	
		$\mu \neq 0$, $m = 1$, $\alpha = 1$	Q_4, Q_5, Q_7	
2.	$f^1 = \lambda u^{-2} v^{-1}$,	$m = 1$, $\alpha = -1$	$e^{\varepsilon t}\left(\partial_t + \varepsilon v\partial_v\right)$, $Q_2,\ Q_3$	11 if $\lambda = 0$
	$f^2 = \sigma u^{-3} + \varepsilon v - \alpha u$	$m = 1$, $\alpha = 1$	$e^{\varepsilon t}\left(\partial_t + \varepsilon v\partial_v\right)$, $Q_4,\ Q_5$	
3.	$f^1 = \lambda v^{\mu+1}$, $f^2 = \sigma v^{\mu-\nu+1}$	$\mu \neq -1$	$(\mu - 2\nu)\, D + \nu t\partial_t$ $-v\partial_v - (\nu+1)u\partial_u$, $\Psi_0(x)\partial_u$	2
4.	$f^1 = \lambda v,\ f^2 = e^{-v}$	$\lambda \neq 0$	$2D - t\partial_t + u\partial_u$ $+\partial_v + \lambda t\partial_u,\ \Psi_0(x)\partial_u$	2
5.	$f^1 = \lambda e^{v},\ f^2 = \sigma e^{v}$	$\lambda\sigma \neq 0$	$D - \partial_v,\ \Psi_0(x)\partial_u$	2
6.	$f^1 = \lambda u^{\nu+1} e^{\mu\frac{v}{u}}$, $f^2 = e^{\mu\frac{v}{u}}(\lambda v + \sigma u)u^{\nu}$	$\mu\lambda \neq 0$	$\mu D - u\partial_v$, $\nu D - u\partial_u - v\partial_v$	
7.	$f^1 = \mu \ln v$, $f^2 = \nu \ln v$	$\nu \neq 0$	$\Psi_0(x)\partial_u$, $D + u\partial_u + v\partial_v$ $+\left(\mu t - \frac{\nu}{2m}x^2\right)\partial_u$	2
8.	$f^1 = 0,\ f^2 = \varepsilon \ln u$	$\varepsilon = \pm 1$	$D - t\partial_t + u\partial_u$ $+\varepsilon t\partial_v,\ t\partial_t + v\partial_v$, $\Phi(u,x)\partial_v$	3, 5, 11
9.	$f^1 = \varepsilon\left(\ln v - \kappa \ln u\right) u$, $f^2 = \varepsilon\left(\ln v - \kappa \ln u\right) v$	$m \neq 2$, $\kappa \neq \frac{m+2}{m-2}$	$(1-\kappa)x_a\partial_{x_a}$ $+2\kappa v\partial_v + 2u\partial_u$, $e^{(1-\kappa)\varepsilon t}\left(u\partial_u + v\partial_v\right)$	1 if $\kappa = 1$
10.	$f^1 = \varepsilon u\left((m+2)\ln u\right.$ $\left.+(2-m)\ln v\right)$, $f^2 = \varepsilon v\left((m+2)\ln u\right.$ $\left.+(2-m)\ln v\right) - \alpha u$	$m \neq 1, 2$ $\alpha = 0$	$Q_1,\ x_a Q_1 - x^2\partial x_a$, $e^{4\varepsilon t}\left(u\partial_u + v\partial_v\right)$	
		$m = 2$, $\alpha = 0$	$H^a\partial_{x_a} - H^a_{x_a} v\partial_v$, $e^{4\varepsilon t}\left(u\partial_u + v\partial_v\right)$	
		$m = 1$, $\alpha = 1$, $\varepsilon = 1$	$Q_2,\ Q_3$, $e^{4t}\left(u\partial_u + v\partial_v\right)$	
		$m = 1$, $\alpha = 1$, $\varepsilon = -1$	$Q_4,\ Q_5$, $e^{-4t}\left(u\partial_u + v\partial_v\right)$	

Table 10. Continued

No	Non-linearities	Conditions	Symmetries	AET (39)
11.	$f^1 = \mu u \ln u,$ $f^2 = \mu v \ln u + \nu v$	$\mu \neq 0$	$e^W \partial_v,$ $e^{\mu t}(u\partial_u + v\partial_v)$	11
12.	$f^1 = \varepsilon v,$ $f^2 = \lambda \frac{v^2}{u} + 2\nu v$ $+\sigma u \ln u$	$\lambda = \pm 1,$ $\sigma = \mp \nu^2$	$Q_8 = e^{\nu t}(\lambda(u\partial_u + v\partial_v)$ $+\nu u \partial_v),\ e^{\nu t} u \partial_v + t Q_8$	
		$\lambda \neq 0,$ $\nu^2 + \lambda\sigma$ $= 1$	$X_\pm = e^{\nu \pm 1}(\lambda(u\partial_u$ $+v\partial_v)$ $+(\nu \pm 1)u\partial_v)$	1 if $\sigma = 0$
		$\lambda \neq 0,$ $\nu^2 + \lambda\sigma$ $= -1$	$e^{\nu t}(\lambda \cos t(u\partial_u + v\partial_v)$ $+(\nu \cos t - \sin t)u\partial_v),$ $e^{\nu t}(\lambda \sin t(u\partial_u + v\partial_v)$ $+(\nu \sin t + \cos t)u\partial_v)$	

The classification of systems of coupled reaction-diffusion equations has been ended.

8 Discussion

We complete the group classification of systems of reaction-diffusion equations started in papers [1], [2], where systems with a diagonal and square diffusion matrix are studied.

The case of triangular diffusion matrix considered in the present paper appears to be rather complicated mainly due to very large number of versions with different symmetries. Indeed, we indicate 54 non-equivalent equations (2) with an invertible triangular diffusion matrix which are presented in Tables 2-6 and 68 equations with a nilpotent diffusion matrix which are present in Tables 8-10 and partly in Tables 2, 3. In addition, 14 classes of equations including first derivatives w.r.t. variables x_ν are collected in Table 7.

In Tables 2, 3 equations defined up to arbitrary functions are presented. In the third column of Table 1 we present the main symmetries of the related equations (2) which do not admit extended symmetries.

In Table 3 we present equations (2) which admit additional equivalence transformations (AET) (39) (indicated in the fourth column) and possible extended symmetries indicated if necessary in the third column after the symbol &.

Tables 4-6 includes the results of classification of equations (2) which are defined up to arbitrary parameters.

The items of Tables 2 and 3 marked by the asterisks are related for equations (2) with $a \neq 0$ and $a = 0$ as well.

Tables 7-10 present the results of group classification of equations (2) with the nilpotent diffusion matrix only.

Note that we did not consider linear equations whose group classification is a rather trivial problem.

Among the classified equations there are only seven of them being invariant w.r.t. the Galilei transformations. In accordance with Item 2 of Table 3 the general form of Galilei-invariant equations (2) is

$$\begin{aligned} &u_t - a\Delta u = uF_1, \\ &v_t - \Delta u - a\Delta v = uF_2 - vF_1 \end{aligned} \tag{50}$$

where F_1 and F_2 are arbitrary functions of variable $\xi = ue^{\frac{v}{u}}$. In Items 3, 5 of Table 4 and Items 1-4 of Table 6 all functions F_1, F_2 are presented which correspond to various possible extensions of the Galilean symmetry. The only system of equations (2) which admits the extended Galilei group including the dilatation and conformal transformations is given by formula (44).

Thus we end the group classification of systems of coupled reaction-diffusion equations (1) started in paper [3] and continued with varying success in [4]-[9] and [1], [2]. The number of non-equivalent equations of this type appears to be enormously large (> 300). Nevertheless, using the approach presented in [1] it was possible to make an effective classification of pairs of coupled reaction-diffusion equations with general diffusion matrix. Moreover, we classify the equations with arbitrary number of independent variables.

We notice that group analysis of reaction-diffusion equations is being intensively developed in many lines including equations with arbitrary elements depending on $t, x, u, u_t, u_x, \cdots$, refer to [10] for a survey. Thus our analysis of systems of diffusion equations is nothing but a part of general study of diffusion models which is currently rather popular.

References

[1] A. G. Nikitin. Group classification of systems of non-linear reaction-diffusion equations with general diffusion matrix. I. Generalised Ginzburg-Landau equations, math-ph/0411027, 2004.

[2] A. G. Nikitin, Group classification of systems of non-linear reaction-diffusion equations with general diffusion matrix. II. Diadonal diffusion matrix, math-ph/0411028, 2004.

[3] Yu. A. Danilov, *Group analysis of the Turing systems and of its analogues*, Preprint of Kurchatov Institute for Atomic Energy IAE-3287/1, 1980.

[4] A. G. Nikitin and R. Wiltshire, in: *Symmetries in Nonlinear Mathematical Physics, Proc. of the Third Int. Conf. , Kiev, July 12-18, 1999, Ed. A.M. Samoilenko* (Inst. of Mathematics of Nat. Acad. Sci. of Ukraine, Kiev, pp. 47-59, 2000).

[5] R. M. Cherniha and J. R. King, J. Phys. A **33**, 267-282, 2000.

[6] R. M. Cherniha and J. R. King, J. Phys. A **33**, 7839-7841, 2000.

[7] A. G. Nikitin and R. Wiltshire, J. Math. Phys. **42** 1667, 2001.

[8] A. G. Nikitin and R. O. Popovych, Ukr. Mat. Zhurn. **53**, 1053-1060, 2001.

[9] R. M. Cherniha and J. R. King, J. Phys. A **36**, 405-425, 2002.

[10] R.O. Popovych and N.M. Ivanova, J. Phys. A **37**, , V.37, 7547-7565, 2004 (see also math-ph/0306035).